DATA LOOK AUGUST 2017 WISCONSIN TOBACCO PREVENTION & CONTROL MOVEMENT

DEATHS

14% of Wisconsin deaths are due to smoking. 2011-2015 Burden of

Tobacco in Wisconsin


EXPOSURE

21%

of Wisconsin high school students are exposed to secondhand smoke in their homes.

2016 WI Youth Tobacco Survey

DISPARITIES PERSIST IN WISCONSIN




of Wisconsin women smoke during pregnancy.

2015 Birth Data; Wisconsin Interactive Statistics on Health

Wisconsin adult smoking rate

2016 Behavioral Risk Factor Surveillance System



2016 WI Youth Tobacco Survey

TRENDS IN SMOKELESS & E-CIGARETTE USE

SMOKELESS

7.4% 0.9%

Wisconsin high school males currently use chewing tobacco, snuff, or dip at a disproportionately higher rate than Wisconsin high school females.

2016 WI Youth Tobacco Survey

ELECTRONIC CIGARETTES

of Wisconsin adults currently use e-cigarettes.

2016 Behavioral Risk Factor Surveillance System

of Wisconsin high school students currently use e-cigarettes.

2016 WI Youth Tobacco Survey 29%

of Wisconsin high school students have ever used e-cigarettes.

2016 WI Youth Tobacco Survey

of Wisconsin high school youth say they would not try e-cigarettes if they were not flavored.

2016 WI Youth Tobacco Survey

THE COST OF TOBACCO



in health care costs are paid in Wisconsin as a result of diseases caused by smoking.

1.6 billion

in lost productivity costs because of illness and premature death from smoking-related illnesses.

2015 Burden of Tobacco in Wisconsin

estimated yearly health care cost of cigarette smoking for every man, woman, and child.

2015 Burden of Tobacco in Wisconsin

